

University of Groningen

Clinical benefit of controversial first line systemic therapies for advanced stage ovarian cancer - ESMO-MCBS scores

Broekman, K. E.; Jalving, M.; van Tinteren, H.; Sessa, C.; Reyners, A. K. L.

Published in:
CANCER TREATMENT REVIEWS

DOI:
[10.1016/j.ctrv.2018.06.008](https://doi.org/10.1016/j.ctrv.2018.06.008)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Broekman, K. E., Jalving, M., van Tinteren, H., Sessa, C., & Reyners, A. K. L. (2018). Clinical benefit of controversial first line systemic therapies for advanced stage ovarian cancer - ESMO-MCBS scores. *CANCER TREATMENT REVIEWS*, 69, 233-242. <https://doi.org/10.1016/j.ctrv.2018.06.008>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Systematic or Meta-analysis Studies

Clinical benefit of controversial first line systemic therapies for advanced stage ovarian cancer – ESMO-MCBS scores

K.E. Broekman^a, M. Jalving^a, H. van Tinteren^b, C. Sessa^c, A.K.L. Reyners^{a,*}^a Department of Medical Oncology, University of Groningen, University Medical Centre Groningen, Hanzeplein 1, Groningen, The Netherlands^b Department of Biometrics, The Netherlands Cancer Institute, Antoni Van Leeuwenhoek Hospital, Plesmanlaan 121, Amsterdam, The Netherlands^c Oncology Institute of Southern Switzerland, Ospedale San Giovanni, CH-6500 Bellinzona, Switzerland

ARTICLE INFO

Keywords:

ESMO-MCBS
Ovarian cancer
Chemotherapy
Targeted therapy
Clinical benefit

ABSTRACT

Background: The magnitude of clinical benefit scale (MCBS) was introduced by the European Society of Medical Oncology (ESMO) to quantify the clinical benefit of therapeutic regimens and to prioritise therapies. It distinguishes curative from palliative treatments and ranks their benefit based on overall survival (OS), progression free survival (PFS), quality of life (QoL) and toxicity. Objective of this study on the first line treatment of ovarian cancer was to evaluate the evidence for the current standard of care using the ESMO-MCBSv1.1 with an emphasis on controversial therapeutic options: intraperitoneal chemotherapy, dose-dense paclitaxel and bevacizumab.

Methods: Phase III trials, published since 1992, investigating first line systemic treatment of Fédération Internationale de Gynécologie et d'Obstétrique (FIGO) stage IIB-IV epithelial ovarian cancer were included. Since most studies included patients with FIGO stage IV disease or incomplete debulking, all treatments were judged to be palliative. Treatments were graded 5 to 1 on the ESMO-MCBSv1.1, where grades 5 and 4 represent a high level of clinical benefit.

Results: 55 studies met the inclusion criteria. ESMO-MCBS scores were calculated for eleven studies that showed a statistically significant benefit of the experimental treatment. Intraperitoneal (ip) cisplatin scored a 4 and 3, but two other studies were negative and therefore not scored on the ESMO-MCBS. Dose-dense paclitaxel showed substantial clinical benefit in one study (score 4), but three studies were negative. Addition of bevacizumab also scored a 4 in one study subgroup including high-risk patients but a 2 in another trial with a larger study population.

Conclusion: Based on ESMO-MCBS scores, dose-dense paclitaxel and intraperitoneal chemotherapy cannot be recommended as standard treatment. Bevacizumab should be considered only in the high-risk population. The ESMO-MCBSv1.1 helps to summarise reported studies on controversial treatment regimens, and identifies their weaknesses.

Introduction

Epithelial ovarian cancer has the highest mortality of all gynaecological cancers [1]. The disease is often diagnosed at a late stage because symptoms only develop once the disease has spread throughout the abdominal cavity. However, even in case of advanced stage disease, long-term survival is possible [2]. First line therapy for advanced disease consists of complete debulking surgery in combination with chemotherapy (carboplatin and paclitaxel given every 3 weeks). Despite extensive research, only few therapies with proven clinical benefit have been added to the therapeutic arsenal during the past decades (Fig. 1). The total costs of ovarian cancer treatment have, however, increased

and vary greatly between European countries, partly due to differences in use of both chemotherapeutic and targeted drugs [3,4]. European Society of Medical Oncology (ESMO) clinical practice guidelines [5] and National Comprehensive Cancer Network (NCCN) guidelines [6] label several first line therapies such as intraperitoneal (ip) cisplatin, dose-dense weekly paclitaxel plus 3-weekly carboplatin, and bevacizumab maintenance therapy as 'optional'. The recent debate on the addition of bevacizumab to first line treatment of advanced ovarian cancer is exemplary for differing policies between European countries with respect to use of new drugs [7].

The inclusion of therapeutic strategies in health insurance plans and health-care outcomes vary significantly between countries [8–12].

* Corresponding author at: Department of Medical Oncology, University Medical Centre Groningen, Hanzeplein 1, Groningen 9713 GZ, The Netherlands.

E-mail address: a.k.l.reyners@umcg.nl (A.K.L. Reyners).

<https://doi.org/10.1016/j.ctrv.2018.06.008>

Received 7 December 2017; Received in revised form 13 June 2018; Accepted 14 June 2018

0305-7372/ © 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Evolution of standard first line systemic therapy in advanced ovarian cancer.

Some therapies provide only a small incremental benefit despite risks of serious toxicity. This makes careful consideration of their value for the quality of life (QoL) of patients and health care costs for society essential. ESMO has developed and validated a tool to assess the magnitude of clinical benefit of cancer medicines: the Magnitude of Clinical Benefit Scale (MCBS) [13,14]. The aim of the ESMO-MCBS is to prioritise the most beneficial treatments and make them available for all patients in Europe. The ESMO-MCBS was designed to rank new drugs based on adequately powered trials, taking into account differences in study design and reported treatment effects. Future anti-cancer drugs or treatments that are approved by the European Medicines Agency (EMA) will be evaluated using the ESMO-MCBS, and interventions showing substantial benefit will be highlighted in the ESMO guidelines. The objective of this study was to re-evaluate the evidence regarding first line treatment of ovarian cancer using the ESMO-MCBSv1.1 with focus on intraperitoneal chemotherapy, dose-dense paclitaxel and bevacizumab.

Methods

Search strategy and scoring

Published, phase III, randomised controlled trials (RCT's) investigating first line systemic treatment of Fédération Internationale de Gynécologie et d'Obstétrique (FIGO) stage IIB-IV ovarian cancer were identified by a PubMed search. The search terms used were: ovarian neoplasms, ovarian cancer, drug therapy, chemotherapy, advanced stage; results were filtered for clinical trials on humans. References of selected articles as well as the Cochrane Library Systematic Reviews and the relevant international guidelines were cross-checked for additional published studies. Studies published from 1992, the introduction of taxanes [15], until March 2018 were included. For studies reporting impending long-term survival data, PubMed was additionally searched specifically for publication of this data. The search was performed independently by two authors (MJ and KEB), in case of different search results these were discussed in the presence of a third investigator (AKLR). For all studies reporting a statistically significant benefit of the experimental treatment over the comparator, the magnitude of clinical benefit was scored by two separate authors (KEB and MJ) on the ESMO-MCBSv1.1. In case of discussion or discrepancy between these two authors' scores, the judgment of a third author was decisive (AKLR). For studies not reporting a statistically significant benefit of the study treatment an ESMO-MCBS score could not be given.

ESMO-MCBS

The ESMO-MCBS [13,14] ranks the value of new cancer therapies in a structured manner, by taking into account reported benefits in terms of longer survival (progression free survival (PFS), disease-free survival

(DFS), overall survival (OS)) and better survival (e.g. QoL, toxicity). To make a comparison possible between trials with different methods, different control groups and endpoints, there are separate scoring forms divided by endpoint and duration of survival of the control group. First, there are separate forms for curative (form 1) and palliative treatments (form 2). Curative or adjuvant treatments are graded A, B or C, with grade A and B corresponding to a substantial clinical benefit. The highest score (A) is given for > 5% improvement of OS at ≥ 3 years, or for improvement of DFS with a hazard ratio (HR) lower limit < 0.65 when mature survival data are lacking. For palliative treatments there are forms for studies with a primary endpoint of OS (form 2a), PFS (form 2b) or QoL, toxicity or response rate (RR) (form 2c). Palliative treatments are graded 5 to 1, where grades 5 and 4 represent a substantial clinical benefit. Recently, a revised version of the ESMO-MCBS was published [14]. In ESMO-MCBSv1.1 a new form 3 was introduced for scoring of single-arm studies in "orphan diseases" and for diseases with "high unmet need" leading to registration of the treatment studied. Furthermore, the threshold for absolute gain in median OS resulting in a score of 5 or 4 was set higher (i.e. more conservative, at 9 months instead of 5 months) for studies with a median survival in the control arm of more than 2 years. The preliminary score for palliative treatments is upgraded when the study treatment shows an improvement in QoL or a reduction in grade 3–4 toxicities impacting on daily well-being. When PFS is the primary endpoint, the preliminary score is downgraded when the study treatment has increased toxicity or does not demonstrate improvement in QoL. All ESMO-MCBS forms can be downloaded online (<http://www.esmo.org/Policy/Magnitude-of-Clinical-Benefit-Scale>).

It can be debated whether treatment of stage IV ovarian cancer, with five-year overall survival of approximately 20%, should be considered curative or palliative. Cure rate depends on the amount of residual disease after surgery. This factor is highly variable, and besides patient and tumour related factors, depends on the surgeon's skills, the location of treatment and the time period in which the study was performed [16]. In this analysis, most studies included patients with stage IV disease and allowed patients with residual masses after debulking surgery to enter. Therefore, all treatments studied in the included trials were labelled 'with palliative intent'.

Validity of included studies

To evaluate the quality and robustness of the included studies and compare studies with and without substantial benefit on the ESMO-MCBS, a Cochrane Institute validity checklist was used [17]. This checklist is a synthesis of the Cochrane Collaboration risk of bias tool [18] and contains ten items that are also included in the CONSORT statement on reporting of RCTs [19] (Supplementary Table 1). It is designed to visualise the validity of studies included in a systematic review, but not to rank them based on a sum score. Weighing of the




Fig. 2. Flow diagram of studies included for analysis according to the ESMO-MCBSv1.1. RCT = randomised controlled trial; HR = hazard ratio.

different items is not incorporated in the checklist; hence there are no threshold values for low or acceptable validity.

Results

The search resulted in 86 hits, of which 55 unique phase III trials met the inclusion criteria. Of these studies, 44 investigated first line therapy and 11 investigated maintenance therapy after first line therapy. There were no single-arm trials leading to registration of the studied treatment in advanced ovarian cancer. QoL data was available for 18 of 55 included studies. Thirteen studies investigating first line treatment and one study on maintenance therapy showed a statistically significant benefit of the experimental treatment. Three of these studies reported a relative risk (RR) only and not a HR, therefore the ESMO-MCBS score could not be calculated. As a result, ESMO-MCBS scoring was performed for eleven of the 55 studies (Fig. 2).

Studies on controversial first line treatments: Intraperitoneal chemotherapy, dose-dense paclitaxel and bevacizumab

Table 1 shows the included studies on intraperitoneal chemotherapy, dose-dense paclitaxel and bevacizumab. Studies showing statistically significant benefit were scored using the ESMO-MCBSv1.1. Ip administration of cisplatin achieved a score of 3 and 4 on ESMO-MCBSv1.1 [20,21]. There were two negative studies on ip treatment [22,23]. Dose dense administration of paclitaxel (80 mg/m² q1w) compared to conventional dose paclitaxel (180 mg/m² q3w) is supported by one study (score 4) [24]. Three negative studies on weekly paclitaxel were identified [25–27]. The addition of bevacizumab to carboplatin and paclitaxel scored a 2 [28] and a 4 [29]. The score of 4 was achieved only in a high-risk population, defined as stage IV disease, inoperable stage III disease, or suboptimally debulked (> 1 cm) stage III disease.

Standard of care options

The current standard first line treatment with carboplatin and paclitaxel is supported by five studies [30–34]. In these five studies carboplatin and paclitaxel were compared to cisplatin with or without cyclophosphamide, the standard of care at that time. ESMO-MCBSv1.1 scores are 4, 4, 4, 3 and 4 respectively. Four of the studies showed

comparable survival but a better QoL [32] or toxicity profile [30,31,33] of the experimental treatment, one study showed an OS benefit [34].

Other studies

Studies investigating other chemotherapeutic agents, anti-hormonal therapy or immune modulating therapies did not show a statistically significant benefit of the experimental treatment over the comparator, nor a better QoL or toxicity profile and are therefore not indicated in the first line treatment of ovarian cancer.

Of the 11 studies investigating maintenance therapy, only pazopanib maintenance treatment showed a statistically significant benefit over the control arm (no maintenance treatment). However, the clinical benefit of maintenance therapy with pazopanib after surgical debulking and first line treatment with carboplatin and paclitaxel could not be qualified as ‘substantial’, with a score of 2 [35].

Table 2 shows the 6 positive studies (5 carboplatin/paclitaxel and 1 pazopanib). All the negative studies on the current standard of care and other treatment regimens are shown in Table 3 [36–71].

The three studies that did not report a HR are summarised in Supplementary Table 2 [72–74]. It concerns one study comparing paclitaxel/cisplatin to cyclophosphamide/cisplatin and two studies investigating ip cisplatin.

Validity of included studies

Validity scores of the 55 included phase III studies are shown in Tables 4a and 4b. Nine studies fulfilled 5 or less of 10 criteria on the Cochrane Institute checklist. Six of these were studies performed before 2000. One of these nine studies showed clinical benefit of carboplatin over cisplatin [31] while the other eight studies did not report clinically meaningful benefits of the experimental treatment. The studies supporting ip administration of cisplatin score 6 out of 10 validity criteria [20,21], as does one study supporting carboplatin over cisplatin added to paclitaxel [33]. All the other studies scored on the ESMO-MCBSv1.1 fulfilled more than 6 validity criteria.

The two studies showing a statistically significant benefit of ip cisplatin but not reporting a HR fulfilled 6/10 validity criteria (Supplementary Table 3). Importantly, ip administration of cisplatin was not the only difference between the control and the intervention arm in these studies. Two doses of carboplatin were added to ip

Table 1
Trials on intraperitoneal chemotherapy, dose-dense paclitaxel and bevacizumab. * if no tumor type specified only EOC patients were included; cm = centimeters; EOC = epithelial ovarian cancer; FTC = fallopian tube cancer; Gr = grade; HR = hazard ratio; ip = intraperitoneal; iv = intravenous; mth = months; ns = not significantly different; OS = overall survival; PFS = progression free survival; PPC = primary peritoneal cancer; QoL = quality of life; RM = residual mass; vs = versus.

Trial name/ first author	Year	Setting	Drug control vs intervention (n)	Primary outcome / design	PFS control	PFS Gain	PFS HR	OS control	OS Gain	OS HR	QoL	Toxicity control vs intervention	ESMO- MCBS v1.1
<i>Intraperitoneal chemotherapy</i>													
Alberts [20]	1996	Stage III after debulking (RM < 2 cm)	Cisplatin iv / cyclophosphamide (n=279) vs cisplatin ip / cyclophosphamide (n=267)	OS				41 mth	8 mth	0.76 (0.61-0.96)		Gr 3-4 neutropenia 69 vs 56% Gr 3-4 abdominal pain 2 vs 18% Gr 3-4 hearing loss 15 vs 5% Gr 3-4 neurotoxicity 25 vs 15%	3
Van Driel [21]	2018	Stage III EOC, FTC, PPC; at least stable disease after 3 cycles of neo- adjuvant treatment	Interval debulking (n=123) vs Interval debulking with HIPEC cisplatin 100mg/m ² (n=122)	PFS OS key secondary				33.9 mth	11.8 mth	0.67 (0.48-0.94)	EORTC QLQ ns	Comparable	4
Kirmani [22]	1994	Stage IIC-IV after debulking	Cisplatin iv / cyclophosphamide iv (n=33) vs cisplatin ip / etoposide ip (n=29)	Complete response rate									
Gadducci [23]	2000	Stage IIC-IV after debulking (RM < 2 cm)	Cisplatin / epidoxorubicin / cyclophosphamide iv (n=57) vs cisplatin ip / epidoxorubicin iv / cyclophosphamide iv (n=54)	OS Crossover ip- iv									
<i>Dose-dense paclitaxel</i>													
IGOC 3016 / Kusumata [24]	2013	Stage II-IV EOC, FTC, PPC	Carboplatin / paclitaxel conventional dose (n=315) vs dose dense (n=312)	PFS OS secondary but data mature				62 mth	38 mth	0.79 (0.63-0.99)		Anemia 44 vs 69% Neuropathy ns	4
Pignatta [25]	2014	Stage Ic-IV after debulking	Carboplatin / paclitaxel conventional dose (n=404) vs dose dense (n=406)	PFS + QoL									
Chan [26]	2016	Stage II-IV EOC, FTC, PPC before / after debulking	Carboplatin / bevacizumab optional / paclitaxel conventional dose (n=346) vs dose dense (n=346)	PFS									
ICON8 [27]	2018	Stage IC high risk – IV EOC, FTC, PPC	Carboplatin / paclitaxel conventional dose vs carboplatin / dose dense paclitaxel vs carboplatin AUC 2 weekly / dose dense paclitaxel	PFS									
<i>Bevacizumab</i>													
GOG 218 / Burger [28]	2011	Stage III or IV EOC, FTC, PPC after debulking	Carboplatin / paclitaxel / placebo (n=625) vs carboplatin / paclitaxel / bevacizumab cycle 2-6 (n=625) vs carboplatin / paclitaxel / bevacizumab cycle 2-22 (n=623)	PFS OS data immature	10.3 mth	0.9 mth bevacizumab cycle 2-6						Gr 2-4 hypertension 7.2 vs 22.9%	2
ICON-7 / Oza [29]	2015	Stage I-IIA (high risk) or stage IIIB-IV EOC, PPC, FTC	Carboplatin / paclitaxel (n=254) vs carboplatin / paclitaxel / bevacizumab (n=248)	OS Prespecified subgroup analysis high risk patients	30.2 mth	9.5 mth >10% OS gain at 3 years 4.4% OS gain at 5 years	0.78 (0.63-0.97) EORTC QLQ ns						4

cisplatin and iv paclitaxel in the experimental arm in one study [72], and in the other study an extra dose of paclitaxel was given per patient in the intervention arm [73]. Two included negative studies (therefore not scored on the ESMO-MCBSv1.1) on ip cisplatin treatment scored low on the validity checklist: 4/10 and 5/10 [22,23].

Discussion

Based on ESMO-MCBS scores, intraperitoneal chemotherapy and dose-dense paclitaxel cannot be recommended as standard treatment. Addition of bevacizumab to first line chemotherapy only showed substantial clinical benefit in a high-risk subgroup of patients (stage IV disease, inoperable stage III disease, or suboptimally debulked (> 1 cm) stage III disease).

Ip chemotherapy using cisplatin does not provide substantial benefit as first line treatment of advanced ovarian cancer according to the score of 3 on the ESMO-MCBSv1.1 [20]. This is the only treatment with a downgraded score (from 4 to 3) on v1.1 of the ESMO-MCBS compared to v1.0. The difference between the two scores is due to the more restrictive criteria for ‘substantial benefit’ on ESMO-MCBSv1.1 (absolute gain ≥ 9 months) compared to version 1.0 (absolute gain ≥ 5 months) [14]. Recently, the results of another study were published [21], showing an improvement of recurrence-free survival (RFS) and OS of the addition of hyperthermic intraperitoneal chemotherapy (HIPEC) with cisplatin 100 mg/m² to interval debulking surgery in stage III ovarian cancer patients not progressing after neo-adjuvant chemotherapy. These outcomes give a score of ‘4’ on the ESMO-MCBS. However, several questions remain, making these results mainly hypothesis-generating and not yet practice-changing at this time [75]. It is unclear whether the extra dose of cisplatin could have caused the observed benefits, and whether hyperthermia is essential for an effect of ip treatment. Furthermore, the results cannot be extrapolated to other populations, such as stage IV patients, patients eligible for primary debulking surgery or patients living in areas with less HIPEC experience. Ip cisplatin is mentioned as optional in guidelines, but it has not been widely adopted. Arguable quality of the available evidence and toxicity are the main arguments against use of ip chemotherapy in an ongoing debate about this treatment modality [76]. OS-results of the large GOG-252 trial comparing dose-dense iv paclitaxel, iv carboplatin and iv bevacizumab (standard arm) to either dose-dense iv paclitaxel, ip carboplatin and iv bevacizumab or iv paclitaxel q3w, ip cisplatin and ip paclitaxel and iv bevacizumab are awaited. However, unfortunately, in this study the treatment arms also differ in more than one way impairing validity [77].

Dose-dense administration of paclitaxel added to carboplatin provides substantial clinical benefit in one study [24]. However, the reported OS benefit could not be confirmed in a recent meta-analysis including three analysed studies (not ICON8, only abstract available) [78], and the three most recent studies on dose-dense paclitaxel showed no PFS or OS-benefit [25–27]. The positive study on dose-dense paclitaxel was conducted in a Japanese patient population. Survival is suggested to be prolonged in Asian patients compared to Caucasian patients [79] potentially explained by genetic differences in drug susceptibility [80]. This might explain the different study results, and this potential bias together with the burden of weekly hospital visits and greater haematological toxicity, means that dose-dense paclitaxel cannot be recommended as part of standard first line treatment.

GOG-218 reports a benefit of adding bevacizumab to first line treatment in ovarian cancer [28], however the score of 2 on the ESMO-MCBSv1.1 does not qualify as ‘substantial improvement’ [13]. In ICON7 an OS benefit for addition of bevacizumab was established in the subgroup of patients at high risk of recurrence only (stage IV disease, inoperable stage III disease, or suboptimally debulked (> 1 cm) stage III disease). This prespecified subgroup analysis scored a ‘4’ [29]. Bevacizumab was approved as a component of first line treatment by the EMA based on the benefit in these studies. However, in several

countries bevacizumab has not been introduced in first line ovarian cancer treatment due to lack of data showing a consistent improvement in OS, concerns about gastro-intestinal toxicity and lack of cost-effectiveness.

The current, post-surgical, standard first line treatment of advanced ovarian cancer, combination chemotherapy with carboplatin and paclitaxel, is strongly supported by four studies that score a substantial clinical benefit on the ESMO-MCBSv1.1 [30–32,34]. There are two studies that did not show clinical benefit of paclitaxel added to platinum compared to platinum monotherapy [41,42]. Possible explanations are cross-over of about 30% from the monotherapy to the combination therapy arm, and a higher dose-intensity of carboplatin in the monotherapy arm of both studies. The largest meta-analysis of different chemotherapeutic schedules in advanced ovarian cancer (60 trials with survival endpoints, N = 15609 women) did show a survival benefit of combination therapy with a platinum and a taxane over platinum monotherapy. This meta-analysis did not only include phase III trials on first line treatment, but also randomised phase II trials and studies on second and third line treatment [81].

Pazopanib maintenance therapy showed clinical benefit based on improvement of PFS. However, due to excess toxicity leading to early treatment discontinuation in nearly one third of the patients, especially in the Asian population, pazopanib did not go into further development in the treatment of ovarian cancer [80,82].

The ESMO-MCBSv1.1 score can only be calculated when there is input of correct data. The first step in using the ESMO-MCBSv1.1 is selection of the most appropriate scoring form, by labelling the goal of the studied treatment as either curative or palliative. All treatments studied in the included trials were labelled ‘with palliative intent’. However, the aforementioned criteria used to label a treatment as ‘with curative’ or ‘palliative intent’ are arbitrary and not evidence based. Seven of the studies included [24,25,33,36,58,72,73] specified multiple primary endpoints but were powered for only one, and five studies [22,45,58,65,74] did not report any power analysis. For one study [41] presented as an equivalence study a non-inferiority design was not reported, therefore scoring on the ESMO-MCBSv1.1 could not be adequately performed. Studies from the past 25 years were included in this analysis. Surgery, but also supportive care including anti-emetics and analgesics have changed over these years. This might influence the comparison of treatment results in this long time period. Current standards for study design are more robust with clear-cut guidelines for reporting results, such as hazard ratios [83]. A recent analysis of 226 contemporary RCTs showed that only one third of modern studies were designed to detect an effect size meeting the ESMO-MCBSv1.0 thresholds [84]. The current analysis includes studies with methodological shortcomings. To take into account the validity of the included studies we used a Cochrane Institute checklist for RCTs. For the six included studies on ip chemotherapy these validity scores clearly illustrate the suboptimal methodological quality of these studies. These study design weaknesses have provoked much of the debate on this treatment strategy. QoL data was only available for 18 of 55 included studies. Palliative studies showing a benefit in PFS but comparable QoL are downgraded one point (form 2B). This makes reporting of QoL data less attractive if no benefit in this endpoint is established. QoL and toxicity are especially important in treatments that are likely palliative, such as in advanced ovarian cancer. Palliative treatments showing non-inferior survival but reduced toxicity can score a ‘4’ on the ESMO-MCBSv1.1 (form 2c). However, reduced toxicity is not clearly defined and the relative weight attributed to different toxicities remains subjective. This is a part of the ESMO-MCBS that could be further elaborated.

Recently, several other methods have been developed to determine the clinical benefit of cancer treatments. The American Society of Clinical Oncology (ASCO) published the ASCO Value Framework, a tool which incorporates clinical benefit, toxicity and also treatment costs [85]. Since the costs of the treatments in the included trials vary between countries, this tool was not used in the current analysis. Another

Table 2
Other positive trials scored for magnitude of clinical benefit on the ESMO-MCBS. * if no tumor type specified only EOC patients were included; cm = centimeters; EOC = epithelial ovarian cancer; FTC = fallopian tube cancer; Gr = grade; HR = hazard ratio; mth = months; OS = overall survival; PFS = progression free survival; PPC = primary peritoneal cancer; QoL = quality of life; RM = residual mass; vs = versus.

Trial name / first author	Year	Setting	Drug control vs intervention (n)	Primary outcome / design	PFS control	PFS Gain	PFS HR	OS control	OS Gain	OS HR	QoL	Toxicity control vs intervention	ESMO-MCBS v1.1
Hannigan [30]	1993	Stage III (RM > 2 cm) or IV after debulking	Cisplatin / cyclophosphamide (n=143) vs carboplatin / cyclophosphamide (n=148)	OS non-inferiority				17 mth	none			Gr 3-4 nausea 17 vs 8% Gr 2-4 renal toxicity 16 vs 2% Gr 2-4 hearing loss 7 vs 0% Gr 3-4 thrombocytopenia 8 vs 24%	4
Taylor [31]	1994	Stage III-IV after debulking	Cisplatin (n=64) vs carboplatin (n=67)	Response rate Crossover	Response duration: 21 mth	none						Crossover due to toxicity 50 vs 3%	4
OVAR-3 / DuBois [32]	2003	Stage IIB-IV after debulking	Cisplatin / paclitaxel (n=386) vs carboplatin / paclitaxel (n=397)	% without progressive disease at 2 years non-inferiority	% without progressive disease at 2 years: 40%	none					EORTC QLQ +13 end of treatment	Gr 3-4 infections 23 vs 36% Gr 3-4 neutropenia 22 vs 36% Gr 3-4 thrombocytopenia 1 vs 13% Gr 3-4 nausea 14 vs 6% Gr 3-4 neuropathy 14 vs 7% Gr 3-4 gastrointestinal 23 vs 10% Gr 3-4 renal 3 vs 1% Gr 3-4 metabolic 8 vs 3% Gr 3-4 thrombocytopenia 5 vs 39%	4
GOG 158 / Osoke [33]	2003	Stage III after debulking (RM < 1 cm)	Cisplatin / paclitaxel (n=400) vs carboplatin / paclitaxel (n=392)	PFS non-inferiority	PFS: 19.1 mth 19.4 mth	none							3
OV-10 / Piccart [34]	2003	Stage IIB-IV after debulking	Cisplatin / cyclophosphamide (n=338) vs cisplatin / paclitaxel (n=324)	OS / % alive at 6.5 years Crossover	OS / % alive at 6.5 years			Approx 36 mth	11% absolute OS gain at 6.5 years	0.75 (0.63-0.90)			4
DuBois [35]	2014	Maintenance stage III-IV EOC, PPC, FTC after debulking and 5 cycles carboplatin / paclitaxel (RM > 2 cm)	Placebo (n=468) vs pazopanib (n=472)	PFS	12.3 mth	5.6 mth	0.77 (0.64-0.91)					Treatment discontinuation 5.6 vs 33.3 %	2

Table 3

Trials showing no statistically significant benefit and therefore not scored on the ESMO-MCBS. * if no tumor type specified only EOC patients were included; # preliminary score of '1' but downgraded for worse QoL; cm = centimeters; EOC = epithelial ovarian cancer; FTC = fallopian tube cancer; ip = intraperitoneal; iv = intravenous; PFS = progression free survival; PPC = primary peritoneal cancer; (p)(c)CR = (pathologic)(clinical)complete response; OS = overall survival; QoL = quality of life; RM = residual mass; vs = versus.

Trial name/first author	Year	Setting*	Drug control vs intervention (n)	Primary outcome
<i>Cisplatin vs carboplatin</i>				
Swenerton [36]	1992	Any stage with macroscopic residual disease after debulking	Cisplatin/cyclophosphamide (n = 210) vs carboplatin/cyclophosphamide (n = 207)	PFS
Meerpohl [37]	1997	Stage III-IV after debulking (RM < 2 cm)	Cisplatin/cyclophosphamide (n = 77) vs carboplatin/cyclophosphamide (n = 81)	PFS
Neijt [38]	2000	Stage IIB-IV after debulking	Cisplatin/paclitaxel (n = 108) vs carboplatin/paclitaxel (n = 100)	PFS
Aravantinos [39]	2005	Stage IIC-IV after debulking	Paclitaxel/carboplatin (n = 121) vs paclitaxel/alternating carboplatin/cisplatin (n = 126)	OS
SCOTROC 4/Banerjee [40]	2013	Stage IC-IV EOC, FTC or PPC after debulking	Carboplatin flat dose (EOC n = 407) vs carboplatin dose escalation (EOC n = 407)	PFS
<i>Paclitaxel</i>				
GOG 132/Muggia [41]	2000	Stage III-IV after debulking (RM > 1 cm)	Cisplatin (n = 200) vs paclitaxel (n = 213) vs cisplatin/paclitaxel (n = 201)	PFS
ICON3 [42]	2002	Any stage after debulking, requiring chemotherapy according to physician	Paclitaxel/carboplatin (n = 478) vs carboplatin (n = 943) and paclitaxel/carboplatin (n = 232) vs cyclophosphamide/doxorubicin/cisplatin (n = 539)	OS
Spriggs [43]	2007	Stage III-IV EOC, FTC, PPC after debulking	Cisplatin/paclitaxel 24 h (n = 140) vs cisplatin/paclitaxel 96 h infusion (n = 140)	PFS
<i>Other cytostatics</i>				
Skarlos [44]	1996	Stage IIC-IV	Carboplatin (n = 73) vs carboplatin/epirubicin/cyclophosphamide (n = 57)	cCR
Wadler [45]	1996	Stage III-IV after debulking or stage I-II with progressive disease or residual disease after irradiation	Melphalan (n = 118) vs cyclophosphamide/hexamethylmelamine/doxorubicin/cisplatin (n = 126)	Clinical response rate
Wils [46]	1999	Stage IC-IV after debulking	Cyclophosphamide/epirubicin/cisplatin (n = 94) vs epirubicin/cisplatin (n = 97)	pCR
SCOTROC 1/Vasey [47]	2004	Stage IC-IV EOC and PPC after debulking	Carboplatin/paclitaxel (n = 538) vs carboplatin/docetaxel (n = 539)	PFS
Du Bois [48]	2006	Stage IIB-IV after debulking	Carboplatin/paclitaxel (n = 635) vs carboplatin/paclitaxel/epirubicin (n = 647)	OS
HIDOC-EIS/Möbus [49]	2007	Stage IIB-IV after debulking	Carboplatin/paclitaxel (n = 71) vs high-dose chemotherapy and peripheral blood stem cell support (n = 78)	PFS
GOG182-ICON 5/Bookman [50]	2009	Stage III-IV EOC and PPC after debulking	Carboplatin/paclitaxel (N = 864) + gemcitabine (n = 864) or liposomal doxorubicin (n = 862) or topotecan (n = 861)	OS
Bolis [51]	2010	Stage III-IV after debulking (RM > 1 cm)	Carboplatin/paclitaxel (n = 170) vs carboplatin/paclitaxel/topotecan (n = 156)	OS
Du Bois [52]	2010	Stage I-IV after debulking	Carboplatin/paclitaxel (n = 882) vs carboplatin/paclitaxel/gemcitabine (n = 862)	OS
Hoskins [53]	2010	Stage IIB-IV EOC, FTC or PPC after debulking	Carboplatin/paclitaxel 8 cycles (n = 410) vs cisplatin/topotecan 4 cycles and carboplatin/paclitaxel 4 cycles (n = 409)	PFS
MITO-2/Pignata [54]	2011	Stage IC-IV after debulking	Carboplatin/paclitaxel (n = 410) vs carboplatin/peg-liposomal doxorubicin (n = 410)	PFS
Fruscio [55]	2011	Stage III-IV after debulking	Cisplatin 3 weekly 6 cycles (n = 139) vs cisplatin weekly 9 cycles (n = 146)	PFS
Lindemann [56]	2012	Stage IIB-IV EOC, PPC or FTC after debulking	Carboplatin/paclitaxel (n = 442) vs carboplatin/paclitaxel/epirubicin (n = 445)	PFS
<i>Antihormonal therapy</i>				
Emons [57]	1996	Stage III-IV	Placebo (n = 66) vs triptorelin (n = 69)	OS
<i>Immunomodulators</i>				
Windbichler [58]	2000	Stage IC-IVa after debulking	Cyclophosphamide/cisplatin (n = 68) vs cyclophosphamide/cisplatin/gamma-interferon (subcutaneous) (n = 65)	PFS
Alberts [59]	2008	Stage III-IV EOC or PPC after debulking	Carboplatin/paclitaxel (n = 421) vs carboplatin/paclitaxel/interferon-gamma (n = 426)	OS
Lhomme [60]	2008	Stage III-IV EOC or PPC after debulking (RM > 1 cm)	Carboplatin/paclitaxel (n = 381) vs carboplatin/paclitaxel/valspodar (n = 381)	PFS
<i>Angiogenesis inhibitors</i>				
AGO-OVAR12/Du Bois [61]#	2016	Stage IIB-IV after debulking	Carboplatin/paclitaxel/placebo (n = 455) vs carboplatin/paclitaxel/nintedanib (n = 911)	PFS
<i>Maintenance therapy</i>				
EORTC 55875/Piccart [62]	2003	Maintenance after debulking and platinum-based chemotherapy (pCR)	Cisplatin ip (n = 76) vs no further treatment (n = 76)	OS
Markman [63]	2003	Maintenance EOC, PPC, FTC after debulking and platinum-based therapy (cCR)	Paclitaxel 3 (n = 128) vs 12 months (n = 134) after complete remission	PFS
MITO-1/De Placido [64]	2004	Maintenance after debulking and 6 cycles carboplatin/paclitaxel	Observation (n = 136) vs topotecan (n = 137)	PFS
Nicoletto [65]	2004	Maintenance after debulking and 5 cycles first line chemotherapy (pCR)	Observation (n = 61) vs 3 courses cisplatin/5-FU (n = 61)	PFS
Hall [66]	2004	Maintenance after debulking and first line chemotherapy	Observation (n = 149) vs interferon-alpha (n = 149)	OS
Alberts [67]	2006	Maintenance stage III after debulking and first line chemotherapy (pCR)	Observation (n = 35) vs interferon-alpha (n = 35)	PFS

(continued on next page)

Table 3 (continued)

Trial name/first author	Year	Setting ^a	Drug control vs intervention (n)	Primary outcome
Hirte [68]	2006	Maintenance stage III-IV after debulking and 6–9 cycles carboplatin/paclitaxel (cCR or RM < 2 cm)	Placebo (n = 121) vs tanomastat (n = 122)	PFS
Pecorelli [69]	2009	Maintenance stage IIB-IV after debulking and 6 cycles paclitaxel/platinum (c/pCR)	Observation (n = 99) vs 6 courses paclitaxel (n = 101)	PFS
Berek [70]	2009	Maintenance stage III-IV after debulking and firstline chemotherapy (cCR)	Placebo (n = 120) vs oregovomab (n = 251)	PFS
MIMOSA/Sabbatini [71]	2013	Maintenance stage III-IV EOC, FTC or PPC after debulking and firstline chemotherapy (cCR)	Placebo (n = 295) vs abagovomab (n = 593)	PFS

Table 4a

Cochrane Institute checklist for validity scores of trials on ip chemotherapy, dose-dense paclitaxel and bevacizumab. Corresponding question Cochrane validity checklist answered with: √ = yes, – = no, ? = unknown (question not answered in referred publication).

Study	Power analysis	Validity items										Total
		1	2	3	4	5	6	7	8	9	10	
<i>Intraperitoneal chemotherapy</i>												
Alberts [20]	√	√	?	–	–	√	√	√	√	√	–	6
van Driel [21]	√	√	?	–	–	√	√	√	–	√	√	6
Kirmani [22]	–	√	√	–	–	√	–	–	–	√	–	4
Gadducci [23]	√	√	√	–	–	–	–	√	√	√	–	5
<i>Dose-dense paclitaxel</i>												
JGOG 3016 [24]	√	√	√	–	–	√	√	√	√	√	√	8
Pignata [25]	√	√	√	–	?	√	√	√	√	√	√	8
Chan [26]	√	√	?	–	?	√	√	√	√	–	–	5
ICON8 [27]	Trial results not published											
<i>Bevacizumab</i>												
GOG 218 [28]	√	√	√	√	√	√	√	√	√	√	–	9
ICON7 2015 [29]	√	√	√	–	–	√	√	√	√	√	–	7

Table 4b

Cochrane Institute checklist for validity scores of trials on other treatments. Corresponding question Cochrane validity checklist answered with: √ = yes, – = no, ? = unknown (question not answered in referred publication).

Study	Power analysis	Validity items										Total
		1	2	3	4	5	6	7	8	9	10	
Hannigan [30]	√	√	√	–	–	√	√	√	√	√	?	7
Taylor [31]	√	√	?	–	–	√	√	?	√	√	?	5
Du Bois [32]	√	√	√	–	–	√	√	√	√	√	–	7
Ozols [33]	√	√	√	–	–	√	√	√	–	√	?	6
Piccart [34]	√	√	√	–	–	√	√	√	√	√	–	7
Du Bois [35]	√	√	√	√	√	√	√	√	√	√	–	9
Swenerton [36]	√	√	√	–	–	√	√	√	√	√	–	7
Meerpohl [37]	√	√	?	–	–	–	√	–	–	√	–	3
Neijt [38]	√	√	?	–	–	√	√	√	√	√	–	6
Aravantinos [39]	√	√	?	–	–	√	–	√	√	√	–	5
SCOTROC4 [40]	√	√	√	–	–	√	√	√	√	√	√	8
GOG 132 [41]	√	√	√	–	–	–	√	√	–	√	√	6
ICON3 [42]	√	√	√	–	–	√	√	√	√	√	√	8
Spriggs [43]	√	√	√	–	–	√	√	√	√	√	–	7
Skarlos [44]	√	√	√	–	–	√	√	√	√	√	–	7
Wadler [45]	–	√	?	–	–	√	√	√	–	√	?	5
Wils [46]	√	√	?	?	?	–	√	√	–	√	–	4
SCOTROC1 [47]	√	√	√	–	–	√	√	√	√	√	–	6
Du Bois 2006 [48]	√	√	√	–	–	√	√	√	√	√	–	7
HIDOC-EIS [49]	√	√	?	–	–	√	√	√	√	√	–	6
ICON5 [50]	√	√	√	–	–	√	√	√	–	√	–	6
Bolis [51]	√	√	√	–	–	√	√	–	√	√	–	6
Du Bois 2010 [52]	√	√	√	–	–	√	√	√	√	√	–	7
Hoskins [53]	√	√	√	–	–	√	√	√	√	√	√	8
MITO-2 [54]	√	√	√	–	–	√	√	√	√	√	–	7
Fruscio [55]	√	√	?	–	–	√	√	√	√	√	?	6
Lindemann [56]	√	√	√	–	–	√	√	√	√	√	–	7
Emons [57]	√	√	√	√	√	√	√	√	–	√	–	8
Windbichler [58]	–	√	√	–	–	√	√	√	√	√	–	7
Alberts 2008 [59]	√	√	√	–	–	√	√	√	√	√	–	7
Lhommé [60]	√	√	√	–	–	√	√	√	–	√	–	6
AGO-OVAR12 [61]	√	√	√	√	√	√	√	√	√	√	–	8
EORTC 55875 [62]	√	√	?	–	–	√	√	√	√	√	–	6
Markman [63]	√	√	?	–	–	√	√	√	√	√	–	6
MITO-1 [64]	√	√	√	–	–	√	√	√	√	√	–	7
Nicoletto [65]	–	√	?	–	–	–	√	√	√	√	–	5
Hall [66]	√	√	√	–	–	√	√	√	√	√	–	7
Alberts 2006 [67]	√	√	√	–	–	√	√	√	√	√	–	7
Hirte [68]	√	√	√	√	?	√	√	√	√	√	–	8
Pecorelli [69]	√	√	√	–	–	√	√	√	√	√	–	7
Berek [70]	√	√	√	√	√	√	√	–	√	√	–	8
MIMOSA [71]	√	√	√	√	√	√	√	√	√	√	–	9

example are the NCCN Evidence Blocks developed to visualise the key measures that form the basis of the recommendations in the NCCN clinical practice guidelines [86].

A major caveat is that ESMO-MCBSv1.1 scores are only calculated for studies showing either a statistically significant survival benefit of the studied treatment, or non-inferior survival but better QoL or less clinically significant grade 3–4 toxicities. Negative studies could be overlooked in treatment advices and guidelines, because of this selection bias. However, keeping this in mind, summarising the clinical benefit of different treatment strategies using the ESMO-MCBS does provide a quick overview of the available body of evidence. A comparable analysis of second and third line treatments could therefore be considered.

Conclusion and future perspectives

Based on ESMO-MCBS scores, dose-dense paclitaxel and intraperitoneal chemotherapy cannot be recommended as standard first line treatment in advanced ovarian cancer. Bevacizumab can be considered in the high-risk population, but is debated because an OS benefit was only shown in this subgroup of patients. The current standard first line treatment in advanced ovarian cancer, combination chemotherapy with carboplatin and paclitaxel, is strongly supported by ranking of the available evidence on the ESMO-MCBSv1.1. ESMO-MCBSv1.1 thresholds for clinical benefit, including QoL analyses and reporting of hazard ratios, should be taken into account in designing future clinical trials.

Funding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest

The authors KE Broekman, M Jalving, H van Tinteren, C Sessa and AKL Reyners declare that they have no competing interests.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.ctrv.2018.06.008>.

References

- [1] Siegel R, Naishadham D, Ahmed J. Cancer statistics, 2013. *CA Cancer J Clin* 2013;63:11–30.
- [2] Dao F, Schlappe BA, Tseng J, et al. Characteristics of 10 year survivors of high-grade serous ovarian carcinoma. *Gynecol Oncol* 2016;141:260–3.
- [3] Kim K, Hermlund E, Hernadi Z, et al. Treatment patterns, health care utilization, and costs of ovarian cancer in central and eastern Europe using a Delphi panel based on a retrospective chart review. *Int J Gynecol Cancer* 2013;23:823–32.
- [4] Greving JP, Vernooij F, Heintz APM, van der Graaf Y, Buskens E. Is centralization of ovarian cancer care warranted? A cost-effectiveness analysis. *Gynecol Oncol* 2009;113:68–74.
- [5] Ledermann JA, Raja FA, Fotopoulou C, Gonzalez-Martin A, Colombo N and Sessa C; ESMO Guidelines Working Group. Newly diagnosed and relapsed epithelial ovarian carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013;24(Suppl. 6):24–32.
- [6] Morgan Jr RJ, Armstrong DK, Alvarez RD, et al. Ovarian Cancer, Version 1.2016, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw* 2016;14:1134–63.
- [7] Hoskins PJ. Time for a level playing field: inequalities in regulatory/approval processes – the example of bevacizumab in epithelial ovarian cancer. *J Clin Oncol* 2015;33:1539–42.
- [8] Gatta G, Trama A, Capocaccia R. Variations in cancer survival and patterns of care across Europe: roles of wealth and health-care organization. *J Natl Cancer Inst Monogr* 2013;46:79–87.
- [9] Luengo-Fernandez R, Leal J, Gray A, Sullivan R. Economic burden of cancer across the European Union: a population-based cost analysis. *Lancet Oncol* 2013;14:1165–74.
- [10] De Angelis R, Sant M, Coleman MP, et al. Cancer survival in Europe 1999–2007 by country and age: results of EUROCARE-5 – a population-based study. *Lancet Oncol* 2014;15:23–34.
- [11] Ades F, Zardavas D, Senterre C, et al. Hurdles and delays in access to anti-cancer drugs in Europe. *Ecancermedicalscience* 2015;8:482.
- [12] Van Harten WH, Wind A, de Paoli P, Saghathian M, Oberst S. Actual costs of cancer drugs in 15 European countries. *Lancet Oncol* 2016;17:18–20.
- [13] Cherny NI, Sullivan R, Dafni U, et al. A standardized, generic, validated approach to stratify the magnitude of clinical benefit that can be anticipated from anti-cancer therapies: the European Society for Medical Oncology Magnitude of Clinical Benefit Scale (ESMO-MCBS). *Ann Oncol* 2015;26:1547–73.
- [14] Cherny NI, Dafni U, Bogaerts J, et al. ESMO-Magnitude of Clinical Benefit Scale version 1.1. *Ann Oncol* 2017; mdx310, <https://doi.org/10.1093/annonc/mdx310>.
- [15] Rowinsky EK, Onetto N, Canetta RM, Arbuck SG. Taxol: the prototypic taxane, an important new class of antitumor agents. *Semin Oncol* 1992;19:646–62.
- [16] Winter WE, Maxwell L, Tian C, et al. Prognostic factors for stage III epithelial ovarian cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2007;25:3621–7.
- [17] Checklist 'Randomised controlled trial' via <http://netherlands.cochrane.org/sites/netherlands.cochrane.org/files/public/uploads/4.4%20Checklist%20Beoordeling%20Randomised%20controlled%20trial%20%28RCT%29.pdf> [accessed 21-08-2017].
- [18] Higgins JP, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- [19] Turner L, Shamseer L, Altman DG, Weeks L, Peters J, Kober T, et al. Consolidated standards of reporting trials (CONSORT) and the completeness of reporting of randomised controlled trials (RCTs) published in medical journals. *Cochrane Database of Systematic Reviews* 2012;(11), Art. No.: MR000030.
- [20] Alberts DS, Liu PY, Hannigan EV, et al. Intraperitoneal cisplatin plus intravenous cyclophosphamide versus intravenous cisplatin plus intravenous cyclophosphamide for stage III ovarian cancer. *New Engl J Med* 1996;335:1950–5.
- [21] Van Driel WJ, Koole SN, Sikorska K, et al. Hyperthermic intraperitoneal chemotherapy in ovarian cancer. *New Engl J Med* 2018;378:230–40.
- [22] Kirmani S, Braly PS, McClay EF, et al. A comparison of intravenous versus intraperitoneal chemotherapy for the initial treatment of ovarian cancer. *Gynecol Oncol* 1994;54:338–44.
- [23] Gadducci A, Carnino F, Chiara S, et al. Intraperitoneal versus intravenous cisplatin in combination with intravenous cyclophosphamide and epidoxorubicin in optimally cytoreduced advanced epithelial ovarian cancer: a randomized trial of the Gruppo Oncologic Nord-Ovest. *Gynecol Oncol* 2000;76:157–62.
- [24] Katsumata N, Yasuda M, Isonishi S, et al. Long-term results of dose-dense paclitaxel and carboplatin versus conventional paclitaxel and carboplatin for treatment of advanced epithelial ovarian, fallopian tube or primary peritoneal cancer (JGOG-3016): a randomised, controlled, open-label trial. *Lancet Oncol* 2013;14:1020–6.
- [25] Pignata S, Scambia G, Katsaros D, et al. Carboplatin plus paclitaxel once a week versus every 3 weeks in patients with advanced ovarian cancer (MITO-7): a randomised, multicentre, open-label, phase 3 trial. *Lancet Oncol* 2014;15:396–405.
- [26] Chan JK, Brady MF, Penson RT, et al. Weekly vs every-3-week paclitaxel and carboplatin for ovarian cancer. *New Engl J Med* 2016;374:738–48.
- [27] Clamp AR, McNeish I, Dean A, et al. ICON 8: A GCG Phase III randomised trial evaluating weekly dose-dense chemotherapy integration in first-line Epithelial Ovarian/Fallopian Tube/Primary Peritoneal Carcinoma (EOC) treatment: Results of Primary Progression-Free Survival (PFS) analysis. Paper presented at: ESMO 2017 Congress; 2017 Sep 8–12; Madrid, Spain.
- [28] Burger RA, Brady MF, Bookman MA, et al. Incorporation of bevacizumab in the primary treatment of ovarian cancer. *New Engl J Med* 2011;365:2473–83.
- [29] Oza AM, Cook AD, Pfisterer J, et al. Standard chemotherapy with or without bevacizumab for women with newly diagnosed ovarian cancer (ICON7): overall survival results of a phase 3 randomised trial. *Lancet Oncol* 2015;16:928–36.
- [30] Hannigan EV, Green S, Alberts DS, et al. Results of a Southwest Oncology Group phase III trial of carboplatin plus cyclophosphamide versus cisplatin plus cyclophosphamide in advanced ovarian cancer. *Oncology* 1993;50(Suppl 2):2–9.
- [31] Taylor AE, Wiltshaw E, Gore ME, Fryatt I, Fisher C. Long-term follow-up of the first randomized study of cisplatin versus carboplatin for advanced epithelial ovarian cancer. *JCO* 1994;12:2066–70.
- [32] Du Bois A, Luck H-J, Meier W, et al. A randomized clinical trial of cisplatin/paclitaxel versus carboplatin/paclitaxel as first-line treatment of ovarian cancer. *J Natl Cancer Institute* 2003;95:1320–30.
- [33] Ozols R, Bundy B, Greer B, et al. Phase III trial of carboplatin and paclitaxel compared with cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2003;17:3194–200.
- [34] Piccart MJ, Bertelsen K, Stuart G, et al. Long-term follow-up confirms a survival advantage of the paclitaxel-cisplatin regimen over the cyclophosphamide-cisplatin combination in advanced ovarian cancer. *Int J Gynecol Cancer* 2003;13(Suppl 2):144–8.
- [35] du Bois A, Floquet A, Kim J-W, et al. Incorporation of pazopanib in maintenance therapy of ovarian cancer. *J Clin Oncol* 2014;32:3374–85.
- [36] Swenerton K, Jeffrey J, Stuart G, et al. Cisplatin-cyclophosphamide versus carboplatin/cyclophosphamide in advanced ovarian cancer: a randomized phase III study of the National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* 1992;10:718–26.
- [37] Meerpohl HG, Sauerbrei W, Kühnle H, Schumacher M, Pfeleiderer A. Randomized study comparing carboplatin/cyclophosphamide and cisplatin/cyclophosphamide as first-line treatment in patients with stage III/IV epithelial ovarian cancer and small volume disease. *Gynecol Oncol* 1997;66:75–84.
- [38] Neijt JP, Engelholm SA, Tuxen MK, et al. Exploratory phase III study of paclitaxel

- and cisplatin versus paclitaxel and carboplatin in advanced ovarian cancer. *J Clin Oncol* 2000;18:3084–92.
- [39] Aravantinos G, Fountzilas G, Kosmidis P, et al. Paclitaxel plus carboplatin versus paclitaxel plus alternating carboplatin and cisplatin for initial treatment of advanced ovarian cancer: long-term efficacy results: a Hellenic Cooperative Oncology Group (HeCOG) study. *Ann Oncol* 2005;16:1116–22.
 - [40] Banerjee S, Rustin G, Paul J, et al. A multicenter, randomized trial of flat dosing versus intrapatient dose escalation of single-agent carboplatin as first-line chemotherapy for advanced ovarian cancer: an SGCTG (SCOTROC 4) and ANZGOG study on behalf of GCG. *Ann Oncol* 2013;24:679–87.
 - [41] Muggia FM, Braly PS, Brady MF, et al. Phase III randomized study of cisplatin versus paclitaxel versus cisplatin and paclitaxel in patients with suboptimal stage III or IV ovarian cancer: a gynecologic oncology group study. *J Clin Oncol* 2000;18:106–15.
 - [42] International Collaborative Ovarian Neoplasms group. Paclitaxel plus carboplatin versus standard chemotherapy with either single-agent carboplatin or cyclophosphamide, doxorubicin and cisplatin in women with ovarian cancer: the ICON3 randomised trial. *Lancet* 2002;360:505–15.
 - [43] Spriggs D, Brady M, Vaccarello L, et al. Phase III randomized trial of intravenous cisplatin plus a 24- or 96-hour infusion of paclitaxel in epithelial ovarian cancer: a Gynecologic Oncology Group study. *J Clin Oncol* 2007;25:4466–71.
 - [44] Skarlos DV, Aravantinos G, Kosmidis P, et al. Carboplatin alone compared with its combination with epirubicin and cyclophosphamide in untreated advanced epithelial ovarian cancer: a Hellenic co-operative oncology group study. *Eur J Cancer* 1996;32A:421–8.
 - [45] Wadler S, Yeap B, Vogl S, Carbone P. Randomized trial of initial therapy with melphalan versus cisplatin-based combination chemotherapy in patients with advanced ovarian carcinoma: initial and long term results – Eastern Cooperative Oncology Group study E2878. *Cancer* 1996;77(733):742.
 - [46] Wils J, van Geuns H, Stoot J, et al. Cyclophosphamide, epirubicin and cisplatin (CEP) versus epirubicin plus cisplatin (EP) in stage Ic-IV ovarian cancer: a randomized phase III trial of the Gynecologic Oncology Group of the Comprehensive Cancer Center Limburg. *Anticancer Drugs* 1999;10:257–61.
 - [47] Vasey PA, Jayson GC, Gordon A, et al. Phase III randomized trial of docetaxel-carboplatin versus paclitaxel-carboplatin as first-line chemotherapy for ovarian carcinoma. *J Natl Cancer Inst* 2004;96:1682–91.
 - [48] Du Bois A, Weber B, Rochon J, et al. Addition of epirubicin as a third drug to carboplatin-paclitaxel in first-line treatment of advanced ovarian cancer: a prospectively randomized gynecologic cancer intergroup trial by the Arbeitsgemeinschaft Gynaekologische Onkologie Ovarian Cancer Study Group and the Groupe d'Investigateurs Nationaux pour l'Etude des Cancers Ovariens. *J Clin Oncol* 2006;24:1127–35.
 - [49] Möbus V, Wandt H, Frickhofen N, et al. Phase III trial of high-dose sequential chemotherapy with peripheral blood stem cell support compared with standard dose chemotherapy for first-line treatment of advanced ovarian cancer: intergroup trial of the AGO-Ovar/AIO and EBMF. *J Clin Oncol* 2007;25:4187–93.
 - [50] Bookman M, Brady M, McGuire W, et al. Evaluation of new platinum-based treatment regimens in advanced-stage ovarian cancer: a Phase III Trial of the Gynecologic Cancer Intergroup. *J Clin Oncol* 2009;27:1419–25.
 - [51] Bolis G, Scarfone G, Raspagliesi F, et al. Paclitaxel/carboplatin versus topotecan/paclitaxel/carboplatin in patients with FIGO suboptimally resected stage III-IV epithelial ovarian cancer: a multicenter, randomized study. *Eur J Cancer* 2010;46:2905–12.
 - [52] Du Bois A, Herrstedt JH, Besard A-C, et al. Phase III trial of carboplatin plus paclitaxel with or without gemcitabine in first-line treatment of epithelial ovarian cancer. *J Clin Oncol* 2010;28:4162–9.
 - [53] Hoskins P, Vergote I, Cervantes A, et al. Advanced ovarian cancer: phase III randomized study of sequential cisplatin-topotecan and carboplatin-paclitaxel vs carboplatin-paclitaxel. *J Natl Cancer Inst* 2010;102:1547–56.
 - [54] Pignata S, Scambia G, Ferrandina G, et al. Carboplatin plus paclitaxel versus carboplatin plus pegylated liposomal doxorubicin as first-line treatment for patients with ovarian cancer: the MITO-2 randomized phase III trial. *J Clin Oncol* 2011;29:3628–35.
 - [55] Frusci R, Garbi A, Parma G, et al. Randomized phase III clinical trial evaluating weekly cisplatin for advanced epithelial ovarian cancer. *J Natl Cancer Inst* 2011;103:347–51.
 - [56] Lindemann K, Christensen RD, Vergote I, et al. First-line treatment of advanced ovarian cancer with paclitaxel/carboplatin with or without epirubicin (TEC versus TC) – a gynecologic cancer intergroup study of the NSGO, EORTC GCG and NCIC CTG. *Ann Oncol* 2012;23:2613–9.
 - [57] Emons G, Ortmann O, Teichert HM, et al. Luteinizing hormone-releasing hormone agonist triptorelin in combination with cytotoxic chemotherapy in patients with advanced ovarian carcinoma. A prospective double blind randomized trial. Decapeptyl Ovarian Cancer Study Group. *Cancer* 1996;78:1452–60.
 - [58] Windbichler GH, Hausmaninger H, Stummvoll W, et al. Interferon-gamma in the first-line therapy of ovarian cancer: a randomized phase III trial. *Br J Cancer* 2000;82:1138–44.
 - [59] Alberts DS, Marth C, Alvarez RD, et al. Randomized phase 3 trial of interferon gamma-1b plus standard carboplatin/paclitaxel versus carboplatin/paclitaxel alone for first-line treatment of advanced ovarian and primary peritoneal carcinomas: Results from a prospectively designed analysis of progression-free survival. *Gynecol Oncol* 2008;109:174–81.
 - [60] Lhomme C, Joly F, Walker J, et al. Phase III study of valsopodar (PSC 833) combined with paclitaxel and carboplatin compared with paclitaxel and carboplatin alone in patients with stage IV or suboptimally debulked stage III epithelial ovarian cancer or primary peritoneal cancer. *J Clin Oncol* 2008;26:2674–82.
 - [61] Du Bois A, Kristensen G, Ray-Coquard I, et al. Standard first-line chemotherapy with or without nintedanib for advanced ovarian cancer (AGO-OVAR 12): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet Oncol* 2016;17:78–89.
 - [62] Piccart MJ, Floquet A, Scarfone G, et al. Intrapertoneal cisplatin versus no further treatment: 8-year results of EORTC 55875, a randomized phase III study in ovarian cancer patients with a pathologically complete remission after platinum-based intravenous chemotherapy. *Int J Gynecol Cancer* 2003;13(suppl 2):196–203.
 - [63] Markman M, Liu PY, Wilczynski S, et al. Phase III randomized trial of 12 versus 3 months of maintenance paclitaxel in patients with advanced ovarian cancer after complete response to platinum and paclitaxel-based chemotherapy: a Southwest Oncology Group and Gynecologic Oncology Group trial. *J Clin Oncol* 2003;21:2460–5.
 - [64] De Placido S, Scambio G, Di Vagno G, et al. Topotecan compared with no therapy after response to surgery and carboplatin/paclitaxel in patients with ovarian cancer: Multicenter Italian Trials in Ovarian Cancer (MITO-1) randomized study. *J Clin Oncol* 2004;22:2635–42.
 - [65] Nicoletti MO, Tumolo S, Falci C, et al. A randomized study of epithelial ovarian cancer: Is chemotherapy useful after complete remission? *Int J Med Sci* 2004;1:116–25.
 - [66] Hall GD, Brown JM, Coleman RE, et al. Maintenance treatment with interferon for advanced ovarian cancer: results from the Northern and Yorkshire Gynaecology Group randomised phase III study. *Br J Cancer* 2004;91:621–6.
 - [67] Alberts DS, Hannigan E, Liu PY, et al. Randomized trial of adjuvant intraperitoneal alpha-interferon in stage III ovarian cancer patients who have no evidence of disease after primary surgery and chemotherapy: An intergroup study. *Gynecol Oncol* 2006;100:133–8.
 - [68] Hirte H, Vergote IB, Jeffrey JR, et al. A phase III randomized trial of BAY 12–9566 (tanomastat) as maintenance therapy in patients with advanced ovarian cancer responsive to primary surgery and paclitaxel/platinum containing chemotherapy: a National Cancer Institute of Canada Clinical Trials Group Study. *Gynecol Oncol* 2006;102:300–8.
 - [69] Pecorelli S, Favalli G, Gadducci A, et al. Phase III trial of observation versus six courses of paclitaxel in patients with advanced epithelial ovarian cancer in complete response after six courses of paclitaxel/platinum-based chemotherapy: final results of the After-6 protocol 1. *J Clin Oncol* 2009;27:4642–8.
 - [70] Berek J, Taylor P, McGuire W, Smith LM, Schultes B, Nicodemus C. Oregovomab maintenance monotherapy does not improve outcomes in advanced ovarian cancer. *J Clin Oncol* 2009;27:418–25.
 - [71] Sabbatini P, Harter P, Scambia G, et al. Abagovomab as maintenance therapy in patients with epithelial ovarian cancer: a phase III trial of the AGO OVAR, COGI, GINECO, and GEICO-the MIMOSA study. *J Clin Oncol* 2013;31:1554–61.
 - [72] Markman M, Bundy BN, Alberts DS, et al. Phase III trial of standard-dose intravenous cisplatin plus paclitaxel versus moderately high-dose carboplatin followed by intravenous paclitaxel and intraperitoneal cisplatin in small-volume stage III ovarian carcinoma: an Intergroup Study of the Gynecologic Oncology Group, Southwestern Oncology Group, and Eastern Cooperative Oncology Group. *J Clin Oncol* 2001;19:1001–7.
 - [73] Armstrong DB, Wenzel B, Huang L, et al. Intrapertoneal cisplatin and paclitaxel in ovarian cancer. *N Engl J Med* 2006;354:34–43.
 - [74] McGuire WP, Hoskins WJ, Brady MF, et al. Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer. *N Engl J Med* 1996;334:1–6.
 - [75] Spriggs DR, Zivanovic O. Ovarian cancer treatment – are we getting warmer? *N Engl J Med* 2018;378:293–4.
 - [76] Bookman MA, Brady MF. Intrapertoneal chemotherapy: long-term outcomes revive a long-running debate. *J Clin Oncol* 2015;33:1424–6.
 - [77] Walker JL, Brady MF, DiSilvestro PA, et al. A phase III trial of bevacizumab with IV versus IP chemotherapy in ovarian, Fallopian tube, and peritoneal carcinoma NCI-supplied agent(s): A GOG/NRG trial (GOG 252). 2016 Society of Gynecologic Oncology Annual Meeting. Late-breaking abstract 6. Presented March 21, 2016.
 - [78] Marchetti C, De Felice F, Musella A, et al. Weekly versus three weeks chemotherapy for advanced ovarian cancer: a meta-analysis. *Oncotarget*. 2016;7:58709–15.
 - [79] Fuh KC, Shin JY, Kapp DS, et al. Survival differences of Asian and Caucasian epithelial ovarian cancer patients in the United States. *Gynecol Oncol* 2015;136:491–7.
 - [80] Oliver KE, McGuire WP. Ovarian cancer and antiangiogenic therapy: caveat emptor. *J Clin Oncol* 2014;32:3353–6.
 - [81] Kyrgiou M, Salanti G, Pavlidis N, Paraskevaidis E, Ioannidis JPA. Survival benefits with diverse chemotherapy regimens for ovarian cancer: meta-analysis of multiple treatments. *J Natl Cancer Inst* 2006;98:1655–63.
 - [82] GlaxoSmithKline: Regulatory update: Votrient (pazopanib) as maintenance therapy for advanced ovarian cancer in the EU. London, United Kingdom, 2014. <http://www.gsk.com/en-gb/media/press-releases/regulatory-update-votrient-pazopanib-as-maintenance-therapy-for-advanced-ovarian-cancer-in-the-eu/> [accessed 21-08-2017].
 - [83] Sobrero AF, Pastorino A, Sargent DJ, Bruzzi P. Raising the bar for antineoplastic agents: how to choose threshold values for superiority trials in advanced solid tumors. *Clin Cancer Res* 2015;21:1036–43.
 - [84] Del Paggio JC, Azariah B, Sullivan R, et al. Do contemporary randomized controlled trials meet ESMO thresholds for meaningful clinical benefit? New cancer therapies and clinical benefit. *Ann Oncol* 2016;28:157–62.
 - [85] Schnipper LE, Davidson NE, Wolins DS, et al. Updating the American Society of Clinical Oncology Value Framework: revisions and reflections in response to comments received. *J Clin Oncol* 2016;34:2925–34.
 - [86] NCCN Evidence Blocks. National Comprehensive Cancer Network. <https://www.nccn.org/evidenceblocks/> [accessed 23-10-2017].